



1/10

FIG. 1






2/10



200710-62934001


3/10



4/10

FIG. 5


#	cDNA
1	1A1
2	2B1 EST
3	2B1
4	2B2
5	3A1 h.
6	3A2
7	4A1
8	$\beta$ -actin
9	GAPDH
10	ext. std.



10043623-011002

5/10

FIG. 6



200710-6294001

6/10

FIG. 7a





FIG. 7b



7/10

FIG. 8a




FIG. 8b




FIG. 8c




FIG. 8d




FIG. 8e






FIG. 8f



DISC-SHAPED PLATFORM

FIG. 8g



MICROTITER / NANOTITER-TYPE  
 DEVICES (96/384/...)

10043625-011002

8/10

MM




FIG. 9c

REGENERATION  
50% UREA




FIG. 9b

PM










FIG. 9a

200110-62951001




9/10

FIG. 10

	epi fluorescence			resonance		
	image	spot means [counts]	signal/ noise*)	image	spot means [counts]	signal/ noise*)
I) proteo-chips incubated with  Cy5-labelled IL6 (500ng/ml, 10 microl, 2h incubation)		46	7.0		1100	69.2
II) Cy5 labelled hCG (50ng/ml, 10 microl, 2h incubation)		32	5.0		646	75.1
III) Cy5-labelled with mixture of anti-IL6 (polyclonal) /IL6 (50ng/ml, 10 microl, 12h incubation)		25	3.8		296	44.1

\*) noise [counts] defined as standard deviation of image background

**Figure 11**



**Figure 12**

